

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/235334898>

Changes in plasma IL4, TNF α and CRP in response to regular passive smoking at home among healthy school children in Khartoum, Sudan

Article in African health sciences · March 2012

Source: PubMed

CITATIONS

3

READS

29

2 authors:



[Tarig Hakim Merghani](#)

University of Khartoum

45 PUBLICATIONS 36 CITATIONS

[SEE PROFILE](#)



[Amal Saeed](#)

University of Khartoum

27 PUBLICATIONS 167 CITATIONS

[SEE PROFILE](#)

Changes in plasma IL4, TNF α and CRP in response to regular passive smoking at home among healthy school children in Khartoum, Sudan

*Merghani TH, Saeed A, Alawad A

Department of Physiology, Faculty of Medicine, University of Khartoum, Sudan

Abstract

Background: Regular exposure to tobacco smoke at home causes airway inflammation and altered cytokine regulation; however, there is variation between individuals of different countries.

Objective: To determine effects of passive smoking on plasma IL4, TNF α , and CRP in healthy male school-children in Khartoum.

Methods: A total of 135 male school-pupils (69 passive smokers and 66 non-smokers) were selected randomly from three primary schools for boys in Khartoum. Information about smoking history was collected from each pupil and his parents using a questionnaire. Plasma CRP, IL4 and TNF α were measured using commercially available ELISA kits.

Results: Plasma IL4 and TNF α in the passive smokers (mean \pm SEM) is at 2.92 ± 0.93 and 19.78 ± 4.67 pg/ml respectively; whereas in the non-smokers. It is at 0.45 ± 0.28 and 5.05 ± 1.54 pg/ml respectively. The difference between the two groups is statistically significant ($p < 0.05$). The mean plasma IL4 and TNF α were lowest with no exposure, higher with moderate exposure, and highest with heavy exposure ($P < 0.05$). Plasma CRP showed insignificant difference between the two groups.

Conclusion: Passive smoking causes significant rise in plasma TNF α and IL4 with a dose dependent effect among school-pupils in Khartoum.

Key words: passive smoking, children, IL4, CRP, TNF α

African Health Sciences 2012; 12(1): 41 - 47

Introduction

Tobacco smoke contains over 4 000 chemicals in the form of particulates and gases.^{1,2} Some of these have marked irritant properties and some are known or suspected carcinogens.¹⁻³ Nicotine which is a weak alkaloid is the most addictive substance.⁴ It is absorbed from the airways to reach the brain in seconds.⁵ It acts on cholinergic receptors in the brain and other sites, causing release of dopamine, endorphins, and stress hormones that exert many effects in the body, including rise in heart rate and blood pressure.^{6,7} Tar, the particulate component of smoke without the water or alkaloid, contains many carcinogens, including polynuclear aromatic hydrocarbons, aromatic amines and N-nitrosamines.⁸ Carbon monoxide binds tightly to haemoglobin in red blood cells, causing reduction in the oxygen-

carrying capacity of red blood cells and therefore predisposes to polycythaemia.⁹ Nitrogen oxides and other oxidants participate in generation of airway inflammation induced by cigarette smoking.¹⁰ The inflammation results in increased mucus production, decreased ciliary movement, and increased mucosal permeability to allergens and therefore increased IgE antibodies & eosinophil count.¹¹

The inflammatory markers C-reactive protein (CRP), interleukin-4 (IL4) & tumor necrosis factor- α "TNF α " play a central role in regulation of inflammatory responses. The CRP is an acute phase glycoprotein produced by the liver. Although its structure is distinct from the immunoglobulins, it shares with them many biological activities. For example, it activates complement,¹² acts as an opsonin¹³ and participates in generation of cytokines that enhance inflammation.¹⁴ Its level rises significantly in inflammation, tissue damage and in the presence of malignant tumors. For this reason, quantitative measurement of CRP is increasingly used as a marker of inflammation and tissue necrosis. Passive smokers who are exposed regularly to environmental tobacco smoke have significantly higher CRP levels in plasma¹⁵⁻¹⁷ indicating an ongoing inflammatory

*Correspondence author:

Dr. Tarig H Merghani
Department of Physiology
Faculty of Medicine
University of Khartoum
P.O.Box 102
Khartoum, Sudan
Email: tarighm@gmail.com

process. However, many studies reported insignificant rise in plasma CRP among passive smokers.^{18,19} It is uncertain whether these variable results are due to genetic factors, environmental factors or related to the short half-life of CRP. The IL4 is a potent multifunctional cytokine. It is produced by T lymphocytes, mast cells and basophils to stimulate growth and differentiation of B lymphocytes that secrete the IgE. Cigarette smoking has been associated with increased serum levels of IL4 and increased risk of developing allergic-like symptoms.²⁰

Many studies reported higher IL4 production by peripheral blood mononuclear cells of smokers, compared to that of non-smokers; However, insignificant findings were also reported. Flouris and his colleagues have recently investigated acute effects of environmental tobacco smoke exposure on plasma cytokines. They reported that IL4 and many other cytokines are highly elevated in response to one hour exposure and remain elevated for at least three hours, but effects of regular long term exposure were not studied.²¹ The TNF α is a potent cytokine that mediates inflammatory and immune reactions.²² It is produced mainly by macrophages to cause inflammation, fever, release of stress hormones, release of acute phase proteins, stimulation of chemotaxis and diapedesis for neutrophils and activation of macrophages.^{23,24} Many studies confirmed that TNF α production and concentrations in body fluids were greater in smokers than in non-smokers.²⁵⁻²⁷ However, several studies reported insignificant results.^{28,29} Recent studies have shown that TNF α plays an important role in the induction of the cigarette related COPD and in the maintenance of airway inflammation.^{30,31} Variation in the results may be explained in part by genetic factor polymorphism.³² That is why certain people are more susceptible to the negative effects of tobacco smoke than others. The aim of this study was to detect changes in plasma level of the inflammatory markers (CRP, IL4 & TNF α) in healthy male school pupils, in response to passive cigarette smoking.

Methods

Study area and population

This is an analytic observational case control study conducted in three governmental primary schools for boys in Khartoum in the year 2009. The schools were selected randomly from 92 primary schools, the total number of public schools for boys in

Khartoum. All pupils in the three schools were approached. Inclusion and exclusion criteria were used to select the test and control groups. The selected pupils were 9 to 14 years old. Parents of the pupils were generally of the average socio-economic class. They lived in houses near the schools of their pupils. Most of the fathers were manual workers or employees and the majority of the mothers were housewives.

Inclusion and exclusion criteria

Inclusion criteria for the test group were: schoolpupil, aged 9 - 14 years, healthy with normal BMI, has no symptoms or signs of acute or chronic medical illness during the last four weeks, no signs of respiratory disease or chest deformity, not on medical treatment, not active smoker, living with at least one smoker at home, exposed regularly to cigarette smoke at home of not less than two cigarettes per day for most days since birth, and not regularly exposed to environmental tobacco smoke or any other type of smoke outside the house.

Inclusion criteria for the control group were: school-pupil, aged 9 - 14 years, healthy with normal BMI, has no symptoms or signs of acute or chronic medical illness during the last four weeks, no signs of respiratory disease or chest deformity, not on medical treatment, not active smoker and not exposed to environmental tobacco smoke or any other type of smoke inside or outside the house.

Exclusion criteria for both groups were: age less than 9 or more than 14 years, abnormal BMI, presence of symptoms of acute or chronic medical illness during the last four weeks, presence of abnormal chest signs or skeletal deformity on clinical examination, being on medical treatment, active smoker, exposed to cigarette smoke, or another type of smoke inside or outside the house (for the non-smokers), or exposed to smoke of less than two cigarettes per day inside the house or to other type of smoke inside or outside the house (for the group of passive smokers). Both groups were matched according to age, sex, height, weight, and area of residence.

Methods of data collection

Questionnaire

Each school pupil was visited at home and interviewed in the presence of his parents to answer questions about his age, health status and smoking habits. Specific questions used for assessment of second-hand smoke exposure included the following

questions: Do you smoke cigarettes? Have you ever smoked cigarettes in the past? Is there anyone in the house who smokes cigarettes? If yes, who? Are you exposed to his cigarette smoke? What is the frequency (number of cigarettes/day)? How many days per week? For how many years? Are you exposed to cigarette smoke outside the house? What is the frequency (number of cigarettes/day)? Are you exposed to any other type of smoke inside or outside the house?). Parents were asked to confirm exposure of their child to cigarette smoke at home and to approve the frequency and duration estimated by their child. The questionnaire paper also requested a signed permission to take a blood sample for measurement of pupil's plasma cytokines.

Evaluation of the degree of exposure

The information obtained from each pupil and his family about frequency of exposure to cigarette smoke at home was used for classification of the pupils into three subgroups: group of no exposure (includes all the non-smokers), group of mild-moderate exposure (exposed to smoke of two - five cigarettes/day) and group of heavy exposure (exposed to smoke of more than five cigarettes/day).

Clinical examination

Each pupil was examined clinically to exclude presence of abnormal clinical signs. Height and weight of each pupil were then measured using standardized scales. The Body mass index (BMI) was calculated for each pupil as weight (in kilograms)/height² (in meters). The BMI results were compared to normal values of BMI, adjusted for age and sex.³³

Preparation and storage of plasma samples

Chemically clean and sterile disposable needles, syringes and swabs were used for all samples. About 5 ml of blood was withdrawn from each pupil from the antecubital vein. The blood was collected in a tube containing EDTA as an anticoagulant. Then the tube was sealed, labelled with the pupil's number, incubated in ice and sent immediately to the laboratory for plasma separation. Plasma was prepared with centrifugation within two hours of venipuncture. The samples were frozen and stored at -20 °C for biochemical analysis.

Biochemical analysis

The inflammatory markers IL4, TNF α were measured using commercially available quantitative

sandwich enzyme-linked immuno-sorbent assay (ELISA) kits according to instructions of the manufacturers. The analyses were performed with 96-well microtiter plate ELISA kits for IL4 & TNF α (from BD Biosciences Pharmingen/USA). CRP was measured with Nycocard Reader II quantitative test (from Axis-Shield/Norway). Test sensitivity was 2 pg/ml for IL4, 2 pg/ml for TNF- α , and 5 mg/L for CRP. Microtiter strips pre-coated with monoclonal antibodies generated against the proteins were used for quantification.

Ethical consideration

The research conforms to the ethical principles of medical research developed by the World Medical Association Declaration of Helsinki.³⁴ Ethical approval was given by the Research Committee (Faculty of Medicine/ University of Khartoum). Approval was obtained from the Unit of School Health/ Ministry of Education. Permission letters were submitted to the headmasters of the selected schools. Written consents was obtained from the parents before entry into the study.

Statistical analysis

Data obtained with questionnaire and biochemical analysis were analyzed using the Statistical Package for the Social Sciences version 16 (SPSS Inc. Chicago, IL, USA). The independent student's t-test was used for comparison between the passive smokers and the non-smokers regarding their physical characteristics (age, weight, height and BMI) and the mean values of CRP, IL4 and TNF α . Statistical significance was accepted when p-value was less than 0.05.

Results

Physical characteristics of pupils in the study group

The mean values of age, weight, height and BMI for the group of passive smokers and the group of non-smokers are presented in table 1. Independent student's t test was used for comparison of the means of the two groups.

Effect of passive smoking on plasma level of inflammatory markers

The mean plasma IL4 and TNF α were significantly higher in the group of passive smokers than the group of non-smokers. The mean plasma CRP was also higher in the passive smokers than the non-smokers; however, the difference was statistically insignificant as shown in table 2.

When the degree of exposure to cigarette smoke was considered, the mean plasma IL4, TNF α , and CRP were lowest with no exposure, higher with mild-moderate degree of exposure, and highest with heavy

exposure. The difference between the three groups was statistically significant for IL4 and TNF α ($p < 0.05$). However, it was insignificant for CRP as shown in table 3.

Table 1: Physical characteristics of pupils in the study group

Parameter	Passive smokers (n = 69)	Non-smokers (n = 66)	P value
	Mean \pm SD	Mean \pm SD	
Age (y)	11.70 \pm 1.34	11.53 \pm 1.2	0.45
Height (cm)	144.24 \pm 9.85	144.00 \pm 8.19	0.87
Weight (Kg)	34.99 \pm 6.83	34.58 \pm 5.69	0.71
BMI	16.66 \pm 1.72	16.60 \pm 1.73	0.83

Differences in physical characteristics between the group passive smokers and the group of non-smokers are not statistically significant

Table 2: Mean plasma levels of inflammatory markers among passive smokers & non-smokers in the study group

Parameter	Passive smokers (n = 69)	Non-smokers (n = 66)
	Mean \pm SEM	Mean \pm SEM
CRP (mg/L)	1.13 \pm 0.34	0.47 \pm 0.25
IL4 (pg/ml)*	2.92 \pm 0.93	0.45 \pm 0.28
TNF α (pg/ml)*	19.78 \pm 4.67	5.05 \pm 1.54

P* < 0.05

The mean plasma IL4 and TNF α are significantly higher in the group of passive smokers than the non-smokers.

Table 3: Plasma levels of inflammatory markers in relation to degree of cigarette smoke exposure (Mean \pm SEM)

Parameter	No exposure (n=66)	Mild-moderate exposure (n=47)	High exposure (n=22)
	Mean \pm SEM	Mean \pm SEM	Mean \pm SEM
cCRP (mg/L)	0.47 \pm 0.25	1.04 \pm 0.36	1.32 \pm 0.72
IL4 (pg/ml)	0.45 \pm 0.28	2.72 \pm 1.24*	3.33 \pm 1.24*
TNF α (pg/ml)	5.05 \pm 1.54	12.42 \pm 3.23*	35.50 \pm 12.47*

P* < 0.05

Mean plasma IL4, TNF α and CRP are lowest with no exposure, higher with mild-moderate degree of exposure, and highest with heavy exposure.

Discussion

Effect of passive smoking on plasma TNF- α

It is well known that cigarette smoke can impair pulmonary immune function and initiate release of inflammatory mediators, hence influencing the development of lung diseases. Recent studies have shown that TNF α plays an important role in the induction of the cigarette-related COPD and in the

maintenance of airway inflammation.³⁵ Experimental studies in rats exposed to tobacco smoke confirmed the rise in TNF α in both serum and broncho-alveolar lavage fluid.^{27,36} In this study, a significant rise in plasma level of TNF α was detected in passive smokers, compared to the control group. This finding indicates that inflammation within the airways of passive

smokers was already established and there would be a risk of progression to COPD with continuous exposure to tobacco smoke. The contribution of passive smoking to the pathogenesis of COPD and its related mortality was not studied in Sudan. However, worldwide, passive smoking is considered an important risk factor for 10-15% of COPD deaths that are not attributed to active smoking.^{37,38} In addition, a large proportion of the active smokers who suffer from COPD have been passive smokers during their childhood.

Effect on plasma IL4

A number of studies proposed that tobacco smoke modifies CD4/CD8 ratio of T cells, increases T helper cells type 2, and increases IL4/IFN α ratio. This is either by increasing IL4 or decreasing IFN α , resulting in activation of B cells to produce IgE antibodies.^{11,20} El-nawawy and colleagues in 1996 conducted a research in Egypt on the effects of passive smoking on frequency of respiratory illnesses, serum IgE and IL4.³⁹ They found higher frequency of respiratory illnesses per year, significantly higher total leucocyte count, higher percentage of eosinophils, and higher serum IgE and IL4 concentrations in pupils of smoking parents compared to the control group. Oryszczynin et al. reported a similar rise in IgE antibodies in adults.⁴⁰ In this study, the rise in plasma level of IL4 in passive smokers was statistically significant. On the other hand, a significant rise in plasma IL4 was detected in the pupils with a high degree of exposure to tobacco smoke compared to the other pupils, again indicating a dose related effect.

Effect on plasma CRP

The acute phase protein CRP is a well studied marker of systemic inflammation⁴¹ and recently has been regarded as a predictor of cardiovascular disease.^{42,43} Although many studies reported significant difference in the level of CRP among passive smokers compared to non-smokers,¹⁵⁻¹⁷ the difference is insignificant in this study. This might be related to the short half-live of CRP or to the lower sensitivity of the quantitative test used for its measurement. However, similar insignificant results were reported in many other studies.

In this study, effects of regular passive smoking at home on plasma inflammatory markers were tested among healthy pupils of the average socio-economic class, selected from three governmental primary schools for boys. Effects on

girls should be investigated to determine whether there is gender difference in cytokine production or not. On the other hand, the effects of passive smoking on affluent pupils of private schools need to be documented because difference in the socio-economic status of pupils in the two types of school may influence the mean values of the results. Previous studies confirmed that high socio-economic factors operating from early life, including the overall household income, affect the risk of developing COPD independently of smoking in both males and females.⁴⁴ On the other hand, low socio-economic status has recently been related to higher levels of inflammatory markers.^{45,46} To avoid variation in the socioeconomic status between passive smokers and non-smokers, pupils of the two groups were selected from the same type of school and matched according to their area of residence. They were also matched according to their age, weight, height and BMI. Those with abnormal BMI were excluded because both obesity and malnutrition cause alterations in the rate of cytokine production.^{47,48} It is worth noting that interviewing pupils in the presence of their parents for assessment of second-hand smoke exposure has the disadvantage because they may not give answers that may make them get criticized by their parents. Nevertheless, it has the advantage that parents can confirm their estimation of frequency and duration of exposure.

Conclusion

These findings confirm that passive smoking causes a significant rise in plasma TNF α and IL4, with a dose dependent effect among school-pupils in Khartoum. Educational programs are highly recommended to increase awareness of parents about the adverse health effects of passive smoking on their children.

References

1. Kiernan VG. Tobacco: a history. London: Hutchison Radius; 1991.
2. Borio G. Tobacco timeline [online]. 1993 [cited 2009 Apr 12]; Available from: [URL:http://www.tobacco.org/tobacco_History.html](http://www.tobacco.org/tobacco_History.html).
3. Doll R. Tobacco: A medical history. *J Urban Health* 1999; 76: 289-313 .
4. The health consequences of smoking: Nicotine addiction: Report of the Advisory Committee of the Surgeon General of the US Public Health Service; 1988.

5. [Gourlay SG, Benowitz NL. Arterio-venous difference in plasma concentration of nicotine and catecholamines and related cardiovascular effects after smoking, nicotine nasal spray and intravenous nicotine. *Clin Pharmacol Ther* 1997; 62: 453-463.](#)
6. [Kapoor D, Jones TH. Smoking and hormones in health and endocrine disorders. *Eur J Endocrinol* 2005; 152 \(4\): 491 -499.](#)
7. [Gidding SS, Xie X, Liu K, Manolio T, Flack J, Perkins L, Gardin J. Smoking has race/gender specific effects on resting cardiac function: the CARDIA study. *Circulation* 1992; 82: 877. Abstract.](#)
8. [Hoffmann D, Djordjevic MV, Hoffmann I. The changing cigarette. *Prev Med* 1997; 26 \(4\): 427 - 434.](#)
9. [Wasserman LR. Cigarette smoking and secondary polycythemia. *JAMA* 1973;224:1654-1657 .](#)
10. [Valença SS, Pimenta WA, Rueff-Barroso CR, Ferreira TS, Resende AC, Moura RS, Porto LC. Involvement of nitric oxide in acute lung inflammation induced by cigarette smoke in the mouse. *Nitric Oxide* 2009;20\(3\):175 -181.](#)
11. [Ronchetti R, Macri F, Ciofetta G, Indinnimeo L, Cutrera R, Bonci E, Antognoni G, Martinez FD. Increased serum IgE and increased prevalence of eosinophilia in 9 year old pupils of smoking parents. *J Allergy Clin Immunol* 1990; 86 \(1\): 400-407.](#)
12. [Kaplan MH, Volanakis JE. Interaction of C-reactive protein complexes with the complement system. Consumption of human complement associated with the reaction of C-reactive protein with pneumococcal C-polysaccharide and with the choline phosphatides lecithin and sphingomyelin. *Journal of immunology* 1974; 112:2135—2147.](#)
13. [Ganrot PO, Kindmark CO. C-reactive protein-a phagocytosis promoting factor. *Scandinavian journal of clinical and laboratory investigation* 1969; 24:215-219.](#)
14. [Du Clos, Terry W. Functions of C- reactive protein. *Ann Med* 2000; 32: 274-278 .](#)
15. [Panagiotakos DB, Pitsavos C, Chrysohoou C, Skoumas J, Masoura C, Toutouzias P, Stefanadis C. Effect of exposure to second hand smoke on inflammation: the ATTICA study. *Am J Med.* 2004; 116: 145–150.](#)
16. [Das I. Raised C-reactive protein levels in serum from smokers. *Clinica Chimica Acta* 1985; 153 :9 –13.](#)
17. [Danesh J, Whincup P, Walker M, Lennon L, Thomson A, Appleby P, Gallimore JR, Pepys MB: Low grade inflammation and coronary heart disease: prospective study and updated meta-analyses. *Brit Med J* 2000; 321:199 – 204.](#)
18. [Helmersson J, Larsson A, Vessby B, Basu S. Active smoking and a history of smoking are associated with enhanced prostaglandin F\(2 \$\alpha\$ \), interleukin-6 and F2-isoprostane formation in elderly men. *Atherosclerosis* 2005;181,201-207 .](#)
19. [Boshtam M, Abbaszadeh M, Rafiei M, Hahparian M, Boshtam M. Comparison of serum levels of CRP and uric acid in active, passive, and non-smokers. *ARYA Journal* 2006;2\(1\):1 -6.](#)
20. [Byron KA, Varigos GA, Wootton AM. IL4 production is increased in cigarette smokers. *Clin Exp Immunol* 1994; 95\(2\): 333 -336.](#)
21. [Flouris AD, Metsios GS, Carrillo AE, Jamurtas AZ, Gourgoulisianis K, Kiropoulos T, Tzatzarakis MN, Tsatsakis AM, Koutedakis Y. Acute and Short-term Effects of Secondhand Smoke on Lung Function and Cytokine Production. *American Journal of Respiratory and Critical Care Medicine* 2009; 179:1029-1033.](#)
22. [Strieter, R., Kunkel, S., Bone, R. Role of Tumor Necrosis Factor-Alpha in Disease States and Inflammation. *Critical Care Medicine* 1993; 21 \(10 Supplement\): S447-S463.](#)
23. [Beutler B, Cerami A. The biology of cachectin/ TNF: A primary host response. *Ann Rev Immunol* 1989;7:625-655 .](#)
24. [Newman I, Wilkinson PJ. Chemotactic activity of lymphotoxin and TNF \$\alpha\$ for human neutrophils. *Immunol* 1989;66:318-320 .](#)
25. [Tappia PS, Troughton KL, Langley-Evans SC, Grimble RF. Cigarette smoking influences cytokine production and antioxidant defenses. *Clin Sci \(London\)* 1995; 88 \(4\): 485 -489.](#)
26. [Kuschner WG, D'Alessandro A, Wong H, Blane PD. Dose-dependent cigarette smoking-related inflammatory responses in healthy adults. *Eur Respir J* 1996;9:1989-1994 .](#)
27. [Pessina GP, Paulesu L, Corradeschi F, Luzzi E, Tanzini M, Aldinucci C, Stefano AD, Bocci V. Chronic cigarette smoking enhances spontaneous release of tumour necrosis factor- \$\alpha\$ from alveolar macrophages of rats. *Mediators Inflamm.* 1993;2\(6\):423 –428.](#)

28. [Keatings VM, Collins PD, Scott DM, Barnes PJ. Differences in interleukin-8 and tumor necrosis factor- \$\alpha\$ in induced sputum from patients with chronic obstructive pulmonary disease or asthma. *Am J Respir Crit Care Med* 1996;153:530-534.](#)
29. [McCrea KA, Ensor JE, Nall K, Bleecker ER, Hasday JD. Altered cytokine regulation in the lungs of cigarette smokers. *Am. J. Respir. Crit. Care Med.*1994;150\(3\):696 -703.](#)
30. [Churg A, Wang RD, Tai H, Wang X, Xie C, Wright JL. Tumor necrosis factor- \$\alpha\$ drives 70% of cigarette smoke-induced emphysema in the mouse. *Am J Respir Crit Care Med* 2004; 170: 492-498.](#)
31. [Churg A, Dai J, Tai H, Xie C, Wright JL: Tumor necrosis factor-alpha is central to acute cigarette smoke-induced inflammation and connective tissue breakdown. *Am J Respir Crit Care Med* 2002, 166:849-854.](#)
32. [Sakao S, Tatsumi K, Igari H, Shino Y, Shirasawa H, Kuriyama T. Association of TNF \$\alpha\$ gene promoter polymorphism with the presence of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001;163:420-422.](#)
33. National Center for Health Statistics, National Center for Chronic Disease Prevention and Health Promotion. Body mass index for age percentiles: Boys, 2 to 20 years. CDC Growth Charts: United states; 2000.
34. World Medical Association Medical Ethics Committee. Updating the WMA Declaration of Helsinki. *Wld Med J* 1999; 45:11-13
35. [Chung KF. Cytokines in chronic obstructive pulmonary disease. *Eur Respir J* 2001;34:Suppl 50s-59s.](#)
36. [Pang B, Wang C, Weng X, Tang X, Zhang H, Niu S, Mao Y, Xin P and Huang X. Lung injury caused by passive smoking and its effects on cytokines in rats. *Zhonghua Yu Fang Yi Xue Za Zhi* 2000; 34 \(2\): 104-105.](#)
37. [Nataraja A. Passive smoking exposure is associated with an increased risk of COPD. *Thorax* 2008;63:48.](#)
38. [Thun MJ, Myers DG, Day-Lally C, Namboodiri MM, Calle EE, Flanders WD, Adams SL, Heath CW. Age and the exposure-response relationships between cigarette smoking and premature death in Cancer Prevention Study II. Changes in cigarette-related disease risks and their implication for prevention and control. Smoking and Tobacco Control, Monograph 8. National Cancer Institute, NIH Publication No. 97-4213., 1997:383-475.](#)
39. [El-Nawawy A, Soliman AT, El-Azzouni O, El-Sayed A, Soheir D, El-Sayed M. Effect of Passive Smoking on Frequency of Respiratory Illnesses and Serum Immunoglobulin-E and Interleukin-4 \(IL4\) Concentrations in Exposed Pupils. *J Trop Pediatr* 1996;42\(3\): 166 -169.](#)
40. [Oryszczyn MP, Annesi-Maesano I, Charpin D, Paty E, Maccario J, Kauffmann F. Relationships of Active and Passive Smoking to Total IgE in Adults of the Epidemiological Study of the Genetics and Environment of Asthma, Bronchial Hyperresponsiveness, and Atopy \(EGEA\). *Am. J. Respir. Crit. Care Med.* 2000;161\(4\):1241-1246.](#)
41. [Young B, Gleeson M, Cripps AW. C-reactive protein: a critical review. *Pathology.* 1991;23:118-24.](#)
42. [Ridker PM, Glynn RJ, Hennekens CH. C-reactive protein adds to the predictive value of total and HDL cholesterol in determining risk of first myocardial infarction. *Circulation* 1998; 97: 2007-2011 .](#)
43. [Ridker PM, Cushman M, Stampfer MJ, Tracy RP, Hennekens CH. Inflammation, aspirin, and risks of cardiovascular disease in apparently healthy men. *N Engl J Med* 1997; 336: 973-979 .](#)
44. [Prescott E, Lange P, Vestbo J. Socioeconomic status, lung function and admission to hospital for COPD: results from the Copenhagen City Heart Study. *Eur Respir J* 1999; 13: 1109-1114 .](#)
45. [Jousilahti P, Salomaa V, Rasi V, Vahtera E, Palosuo T. Association of markers of systemic inflammation, C reactive protein, serum amyloid A, and fibrinogen, with socioeconomic status. *J Epidemiol Community Health.* 2003;57:730-733.](#)
46. [Owen N, Poulton T, Hay FC, Mohamed-Ali V, Steptoe A. Socioeconomic status, C-reactive protein, immune factors, and responses to acute mental stress. *Brain Behav Immun.* 2003;17:286-295 .](#)
47. [Popko K, Gorska E, Stelmaszczyk-Emmel A, Plywaczewski R, Stoklosa A, Gorecka D, Pyrzak B,](#)
48. [González-Martínez H, Rodríguez L, Nájera O, Cruz D, Miliar A, Domínguez A, Sánchez F, Graniel J, González-Torres MC. Expression of Cytokine mRNA in Lymphocytes of Malnourished Children. *J of Clin Immunol* 2008;28\(5\):593-599.](#)